arXiv:1112.6158v1 [physics.atom-ph] 28 Dec 2011

Confining diffuse potential versus square-well-potential in modeling A@C$_{60}$ atoms

V. K. Dolmatov, J. L. King, and J. C. Oglesby
Department of Physics and Earth Science, University of North Alabama, Florence, Alabama 35632, USA

(Dated: December 30, 2011)

It is shown that discontinuity, inherit to a square-well potential which is often used for mimicking the C$_{60}$ cage potential, results neither in qualitative nor, what is even more important, quantitative artifacts in problems of endohedral A@C$_{60}$ atoms, such as their photoionization. Moreover, it is demonstrated that, upon mimicking a square-well potential by a potential with diffuse boarders, calculated photoionization spectra are largely insensitive to the degree η of diffuseness of the potential boarders, in a reasonably broad range of η's. The proof is based on results of comparison between calculated data for H@C$_{60}$ and Xe@C$_{60}$ photoionization cross sections and photoelectron angular asymmetries obtained by modeling the C$_{60}$ cage by confining square-well and diffuse potentials. Thus, it matters little which of these potentials to use in A@C$_{60}$ problems. The diffuse potential is modeled by a combination of two Woods-Saxon potentials. Calculated data for the photoionization of Xe@C$_{60}$ are separately detailed as well, since they are important stand-alone qualities.

PACS numbers: PACS: 32.80.Fb, 32.30.-r, 31.15.V-

I. INTRODUCTION

Photoionization of atoms A encapsulated inside the hollow interior of the C$_{60}$ fullerene cage, labeled as A@C$_{60}$ and referred to as endohedral or confined atoms, has been an *ad hoc* topic of numerous theoretical (see review papers [1, 2] (and references therein in addition to other references in the present paper) and experimental [3] (and references therein) studies in recent years. An *ab initio*, based on first principles theory of A@C$_{60}$ photoionization has not been developed yet, in view of a formidable complexity of the problem. Perhaps, the most sophisticated theory here is represented by a time-dependent local density approximation (TDLDA), see, e.g., Ref. [4] and references therein. However, TDLDA theory has obvious drawbacks in view of an unsatisfactory agreement with the existing experimental data on the Xe@C$_{60}^{+}$ 4d photoionization [3]. Meanwhile, many important insights into the problem can be, and have been, unraveled on the basis of simpler empirical models based on the modeling of C$_{60}$ confinement by pseudo-potentials, such as a δ-function-like potential [5, 6] (and references therein) or square-well-potential, $U_{\rm SWP}(r)$. The latter is defined as

$$U_{\rm SWP}(r) = \begin{cases} -U_0, & \text{if } R_0 \le r \le R_0 + \Delta \\ 0 & \text{otherwise.} \end{cases} \tag{1}$$

Here, R_0 is the inner radius of C$_{60}$, Δ is the thickness of the C$_{60}$ wall, and U_0 is the potential depth. The square-well-potential modeling of C$_{60}$ has become quite popular among various researchers and been used on numerous occasions in the field of C$_{60}$ and A@C$_{60}$ theoretical studies, see, e.g., Refs. [1, 2, 7–19] and references therein.

The potential $U_{\rm SWP}(r)$, however, is discontinuous at its boarders. A possible emergence of qualitative and, especially, quantitative artifacts, associated with said discontinuity, in calculated photoionization cross sections of, and photoelectron angular-asymmetries from, A@C$_{60}$ atoms has not been detailed in literature except for a couple of brief remarks dropped on the subject in passing in Refs. [20, 21]. In particular, it is because of the lack of such knowledge that some authors, see, e.g., Ref. [22], have recently proposed to discard the square-well-potential modeling of the C$_{60}$ cage on grounds of discontinuity of the potential, and replace it by a Gaussian-function-like model potential. This proposition, we argue, is wrong. The proposed potential continuously changes strongly everywhere inside the C$_{60}$ cage, thereby having no compact boarders. This contradicts a recent Fourier imaging study of the experimental C$_{60}$ photoionization cross section [23]. From the study, the fact that the C$_{60}$ has well-defined, sharp boarders is obvious. In short, following the logic line of Ref. [20, 23], photoionization of an atomic cluster occurs with the greatest probability where the cluster's potential changes sharply. This is obvious from the acceleration form gage for a dipole photoionization amplitude, D, namely, $D \propto (\psi_f|\Delta_r V(r)|\psi_i)$. On grounds of this, the Fourier imaging study [23] of the experimental C$_{60}$ photoionization cross section [3] has proven that spherical boundaries of C$_{60}$ are rather sharp. Hence, the C$_{60}$ potential should be reminiscent of a square-well potential $U_{\rm SWP}(r)$, to a good approximation, rather than a soft Gaussian-function-like potential peaked at the C$_{60}$ mid-wall.

Surely, a C$_{60}$ confining potential with diffuse (but well defined) boarders is much more realistic than a square-well-potential with infinitely sharp edges. Correspondingly, replacement of the latter with the former in A@C$_{60}$ problems would be, of course, worthwhile, for the sake of consistency. With this in mind, it is not at all clear *apriori* to what degree this may qualitatively and quantitatively affect/alter results of numerous previous calculational predictions on A@C$_{60}$ photoionization made on the basis of the square-well-potential concept, and whether such replacement will improve agreement with experiment or make it worse. The need for clarifying this issue for a deeper insight into the structure and spectra of A@C$_{60}$ atoms is additionally signified by the fact that the square-well-potential model of A@C$_{60}$ atoms has been used abundantly over the years which has resulted in a large array of predicted data and phenomena that might need to be re-studied/re-calculated with an eye on more realistic diffuse potential boarders.

It is the aim of this paper to dot i's and cross t's concerning possible consequences of replacement of the confining square-well-potential $U_{\rm SWP}(r)$ by a diffuse potential $U_{\rm DP}(r)$ in A@C$_{60}$ problems. To achieve this end, we calculate the photoionization cross sections of, and photoelectron angular distribution from, inner and valence subshells of A@C$_{60}$ atoms by utilizing both $U_{\rm SWP}(r)$ and $U_{\rm DP}(r)$ [which, nevertheless, is reminiscent of $U_{\rm SWP}(r)$, see next section] as model pseudo-potentials that mimic the C$_{60}$ potential. It is proven in the present work that discontinuity of the $U_{\rm SWP}(r)$ potential in question does not matter at all, in photoionization problems. It leads neither to qualitative nor quantitative artifacts in calculated photoionization characteristics of A@C$_{60}$ atoms compared to calculated data obtained with the use of the diffuse potential $U_{\rm DP}(r)$. Moreover, it is found that the degree of diffuseness of the $U_{\rm DP}(r)$ potential matters surprisingly little - a fraction-to-nothing - in a relatively broad range of its values.

Atomic units (a.u.) are used throughout this paper.

II. THEORY, RESULTS, AND DISCUSSION

A. Review of theory

In this work, to model the C$_{60}$ cage by a square-well-potential we use $U_{\rm SWP}(r)$ given by Eq. (1).

To mimic the C$_{60}$ cage by a diffuse potential $U_{\rm DP}(r)$, which is somewhat reminiscent of $U_{\rm SWP}(r)$, we form the latter by the appropriately defined combination of two Woods-Saxon potentials:

$$U_{\rm DP}(r) = \left.\frac{2U_0}{1+\exp(\frac{R_0-r}{\eta})}\right|_{r\le R_0+\frac{1}{2}\Delta} + \left.\frac{2U_0}{1+\exp(\frac{r-R_0-\Delta}{\eta})}\right|_{r>R_0+\frac{1}{2}\Delta}. \qquad (2)$$

Here, η is the diffuseness parameter, and R_0, U_0, and Δ are the square-well-potential parameters.

Concerning photoionization of A@C$_{60}$ atoms, we focus on the photoionization cross section $\sigma_{n\ell}(\omega)$ of a $n\ell$-subshell of A@C$_{60}$ as well as dipole photoelectron angular-asymmetry parameter $\beta_{n\ell}(\omega)$ which governs the pattern of a $n\ell$-photoelectron angular distribution, see, e.g., Refs. [24, 25] for details. Expressions for $\sigma_{n\ell}(\omega)$ and $\beta_{n\ell}(\omega)$ for A@C$_{60}$ are the same as for free atoms presented in the above references. We write,

$$\sigma_{n\ell} = \frac{4\pi^2\alpha N_{n\ell}}{3(2\ell+1)}\omega[\ell|D_{\ell-1}|^2 + (\ell+1)|D_{\ell+1}|^2], \qquad (3)$$

and we recast $\beta_{n\ell}(\omega)$ as

$$\beta_{n\ell} = \frac{\ell(\ell-1)\rho^2 - 6\ell(\ell+1)\rho\cos\Phi + (\ell+1)(\ell+2)}{(2\ell+1)(\rho^2\ell+\ell+1)}, \qquad (4)$$

where

$$\rho = \frac{|D_{\ell-1}|}{|D_{\ell+1}|}, \quad \Phi = \delta_{\ell+1} - \delta_{\ell-1}. \qquad (5)$$

In the above equations, ω is the photon energy, $N_{n\ell}$ is the number of electrons in a $n\ell$ subshell, α is the fine-structure constant, $D_{\ell\pm1}$ are radial dipole photoionization amplitudes, and $\delta_{\ell\pm1}$ are phase shifts of $D_{\ell\pm1}$. Note that the quantities $\sigma_{n\ell}$, $\beta_{n\ell}$, $D_{\ell\pm1}$, ρ, $\delta_{l\pm1}$, and Φ all depend upon photon energy ω; the explicit dependence is omitted in the above equations for reasons of simplicity.

In a one-electron approximation, $D_{\ell\pm1}$ is defined as

$$D_{\ell\pm1} = \int_0^\infty P_{\epsilon\ell\pm1}(r) r P_{n\ell}(r) dr, \qquad (6)$$

where $P_{n\ell}(r)$ and $P_{\epsilon\ell}(r)$ are one-electron radial wavefunctions of the bound and continuous states, respectively. In the present work, these wavefunctions and energies $E_{n\ell}$ of a discrete spectrum are the solutions of a radial Hartree-Fock (HF) equation accounting for a $U_{\rm C_{60}}(r)$ confinement:

$$[\hat{H}^r + U_{\rm C_{60}}(r)]P_{n\ell}(r) = E_{n(\epsilon)\ell}P_{n(\epsilon)\ell}(r). \qquad (7)$$

Here, $\hat{H}^r$ is a radial part of the HF Hamiltonian which is identical to that for a free atom, see, e.g., Ref. [24] for the latter, and $U_{\rm C_{60}}(r)$ is either the square-well-potential $U_{\rm SWP}(r)$, Eq. (1), or diffuse potential $U_{\rm DP}(r)$, Eq. (2). In a particular case of an endohedral single-electron hydrogen atom, H@C$_{60}$, Eq. (7) reduces to a radial Schrödinger equation in the presence of the C$_{60}$ confinement:

$$-\frac{1}{2}\frac{d^2P_{n(\epsilon)\ell}}{dr^2} + \left[\frac{-1}{r} + \frac{\ell(\ell+1)}{2r^2} + U_{\rm C_{60}}(r)\right]P_{n(\epsilon)\ell}(r) = E_{n(\epsilon)\ell}P_{n(\epsilon)\ell}(r). \qquad (8)$$

To calculate $D_{\ell\pm1}$ beyond a one-electron HF approximation, i.e., to account for impacts of electron correlation on the photoionization process, we utilize a random phase approximation with exchange (RPAE) [24]; the reader is referred to Ref. [24] for details of the RPAE theory.

B. Results and discussion

In this work, when approximating the C$_{60}$ potential by a square-well or diffuse potential, we use the proposed in Ref. [26] values of $U_0 = -0.422$, $\Delta = 1.25$, and $R_0 = 6.01$ a.u. rather than the earlier used $U_0 = -0.302$, $\Delta = 1.9$, and $R_0 = 5.89$ a.u., see, e.g., Refs. [1, 2, 13]. This is because the new parameters were shown [26] to result in a much better match to experimental photoionization



FIG. 1: (Color online) The square-well-potential $U_{\rm SWP}(r)$, Eq. (1), with $U_0 = -0.422$, $\Delta = 1.25$, and $R_0 = 6.01$ a.u. (dotted line), as well as the diffuse potential $U_{\rm DP}(r)$, Eq. (2), with the diffuseness parameter $\eta = 0.01$ (solid line) and $\eta = 0.1$ (dashed line). See text for details.



FIG. 2: (Color online) Hydrogen $1s$ photoionization cross section of H@C$_{60}$ calculated with the use of the square-well-potential $U_{\rm SWP}(r)$ (open circles), diffuse potential $U_{\rm DP}(r)$ with $\eta = 0.01$ (solid line) and $\eta = 0.1$ (dashed line), and of free hydrogen, as marked.

spectra of endohedral atoms (of Xe@C$_{60}^{+}$ [3], in particular).

Calculated data for the thus defined potentials $U_{\rm SWP}(r)$ and $U_{\rm DP}(r)$ are compared with each other in Fig. 1 for two different values of the diffuseness parameter η of $U_{\rm DP}(r)$, namely, $\eta = 0.01$ and 0.1 . One can see from Fig. 1 that $U_{\rm DP}(r)$ with $\eta = 0.01$ matches the square-well-potential practically precisely. On the contrary, the substitution of $\eta = 0.1$ in $U_{\rm DP}(r)$ results both in relatively strongly diffuse boarders and bottom of $U_{\rm DP}(r)$, so that the latter noticeably deviates from the square-well-potential.

In the following, we detail results of comparison between $\sigma_{n\ell}(\omega)$ and $\beta_{n\ell}(\omega)$ calculated in the presence of the above determined $U_{\rm SWP}(r)$ and $U_{\rm DP}(r)$ confinement.

As the first step, in order to avoid various possible complications associated with electron correlation in multielectron atoms, let us discuss a "clean" case - the photoionization of the $1s$ ground-state of the endohedral hydrogen atom, H@C$_{60}$. Corresponding results are depicted in Fig. 2. First, we note that the photoionization cross section $\sigma_{1s}(\omega)$ of H@C$_{60}$ calculated with the use of the square-well-potential $U_{\rm SWP}(r)$ and diffuse potential $U_{\rm DP}(r)$ with a small diffuseness parameter $\eta = 0.01$ are undistinguishable from each. Hence, discontinuity of the square-well-potential does not matter at all in this problem. Second, what comes as a big surprise, however, is that calculated data obtained with the use of a much greater diffuseness parameter, $\eta = 0.1$, for which $U_{\rm DP}(r)$ deviates from a square-well-potential considerably (see Fig. 1), are practically the same as calculated results obtained with $\eta = 0.01$ or with the use of the square-well-potential. This is the initial indication that the photoionization cross section of an endohedral atom may be rather insensitive to the degree of diffuseness of the confining potential boarders, in reasonably large limits. As a note on an independent issue, one can see that the photoionization cross section of free hydrogen differs from that of H@C$_{60}$ by the presence of a very well developed maximum at about 25 eV and another less developed maximum at about 43 eV in the latter. These are called confinement resonances, i.e., resonances that occur due to a constructive interference of the outgoing photoelectron wave and photoelectron waves scattered off the C$_{60}$ cage [1, 2, 5] (and references therein). Confinement resonances have only recently been experimentally proven as reality [3].

In the next, we investigate whether the above findings are valid for photoionization of inner-shells and valence-shells of endohedral multielectron atoms A@C$_{60}$ as well. To date, only photoionization of endohedral Xe@C$_{60}^{+}$ has been reliably measured experimentally [3]. Therefore, we choose photoionization of Xe@C$_{60}$ as the case study to learn how discontinuity of $U_{\rm SWP}(r)$ and diffuseness of $U_{\rm DP}(r)$ may affect photoionization cross sections $\sigma_{n\ell}(\omega)$ of, and photoelectron angular-asymmetry parameters $\beta_{n\ell}(\omega)$ from, the Xe@C$_{60}$ inner $4d^{10}$ and valence $5p^6$ subshells. Note, $\beta_{n\ell}(\omega)$ depends on phase shifts $\delta_{\ell\pm1}$



FIG. 3: (Color online) RPAE calculated data for $\sigma_{4d}(\omega)$ and $\beta_{4d}(\omega)$ of Xe@C$_{60}$ obtained with the use of the square-well-potential $U_{\rm SWP}(r)$ (dashed lines) and diffuse potential $U_{\rm DP}(r)$ with $\eta = 0.1$ (solid lines), as well as corresponding HF calculated data obtained with the use of $U_{\rm DP}(r)$ with $\eta = 0.1$ (dotted lines), as marked. Also depicted are corresponding RPAE calculated data for free Xe (dashed-dotted lines) and experimental data for $\sigma_{4d}(\omega)$ of Xe@C$_{60}^{+}$ [3]. Experimental data were shifted by 4.2 eV towards higher photon energies as well as multiplied by 10 to compare with theory (see discussion in Appendix).

of photoionization matrix elements. Hence, corresponding calculated data will provide, although implicitly, the information on sensitivity of phase shifts $\delta_{\ell\pm1}$ to diffuseness of a confining potential as well. RPAE calculated data for $\sigma_{4d}(\omega)$ and $\beta_{4d}(\omega)$, as well as for $\sigma_{5p}(\omega)$ and $\beta_{5p}(\omega)$, are depicted in Fig. 3 and Fig. 4, respectively.

One can see from Figs. 3 and 4 (compare dashed curves marked as "$U_{\rm SWP}$" with solid curves marked as "$U_{\rm DP}$, $\eta = 0.1$") that, similar to the H@C$_{60}$ case, there are little-to-no differences both in calculated $\sigma_{n\ell}(\omega)$'s and $\beta_{n\ell}(\omega)$'s obtained with accounting for a square-well-potential confinement versus those obtained with a diffuse confinement potential even with as large degree of diffuseness as $\eta = 0.1$. Thus, photoionization spectra both of a single-electron and multielectron endohedral atom A@C$_{60}$ are equally insensitive to discontinuity in $U_{\rm SWP}$ or diffuseness of $U_{\rm DP}$; the implication is that the findings carry the general meaning.



FIG. 4: (Color online) RPAE calculated data for Xe@C$_{60}$ 5p photoionization cross section, $\sigma_{5p}(\omega)$, and dipole photoelectron angular-asymmetry parameter, $\beta_{5p}(\omega)$, calculated with the use of the square-well-potential $U_{\rm SWP}(r)$ (dashed lines) and diffuse potential $U_{\rm DP}(r)$ with $\eta = 0.1$ (solid lines). Also depicted are RPAE calculated data for $\sigma_{5p}(\omega)$ and $\beta_{5p}(\omega)$ for free Xe (dashed-dotted lines).

III. CONCLUSION

To summarize, it has been proven in this paper that discontinuity of a square-well-potential does not lead to any artifacts, whether quantitative or qualitative, in photoionization spectra of endohedral atoms. The square-well-potential model of C$_{60}$ is as good as its modeling by a diffuse potential. Both of these potentials lead to practically identical calculated data for photoionization spectra. The latter appear to be largely insensitive to the degree of diffuseness. Hence, either of the two potentials is equally suitable for mimicking the C$_{60}$ cage. In particular, switching from a confining square-well-potential to a more realistic diffuse potential, if wanted, will not require a re-study of a rich variety of important results on A@C$_{60}$ ionization phenomena obtained to date on the basis of a confining square-well-potential concept.

Finally (see Appendix), since the values of square-well-potential parameters $R_0 = 6.01$, $\Delta = 1.25$, and $U_0 = 0.422$ a.u., empirically deduced in Ref. [26] and exploited in the present work, lead to a reasonable agreement with experiment (see Fig. 3), we suggest the users of the square-well-potential modeling of A@C$_{60}$ to utilize these updated parameters in their future work on the subject.

Acknowledgments

This work was supported by NSF Grant No. PHY-0969386. Dr. S. T. Manson is thanked for bringing the Ref. [20] to the authors' attention.

Appendix

To finalize the paper, we now briefly comment on particular features and structures in the calculated spectra of Xe@C_{60} depicted in Figs. 3 and 4, since they are important stand-alone qualities.

1. The $4d$ photoionization of Xe@C_{60}

To start with, the three strong oscillations in $\sigma_{4d}(\omega)$ of Xe@C_{60} about $\sigma_{4d}(\omega)$ of free Xe are confinement resonances (see the definition of confinement resonances given above in this paper). They were previously theoretically studied and detailed in Refs. [4, 5, 13, 15, 26] and experimentally in Ref. [3]. Results of the measurement are by about the factor of 10 smaller than the cited theoretical data. This is because the measured channel of the Xe@C_{60}^{+} $4d$ photoionization might [3] account for only about 10% of the Xe $4d$ oscillator. For this reason the experimental data depicted in Fig. 3 of the present paper are multiplied by 10, to ease the comparison with theory. One can see that calculated results for $\sigma_{4d}(\omega)$ of Xe@C_{60} obtained in the present work with the square-well-parameters $R_0 = 6.01$, $\Delta = 1.25$, and $U_0 = 0.422$ a.u. are in a reasonable agreement with main structures of the experimental spectrum, to a good approximation. Interesting, HF calculated data for $\sigma_{4d}(\omega)$ of Xe@C_{60} seem to be even in a better, than RPAE data, agreement with experiment beyond the photon energy of about 100 eV. Note, the comparison of our calculated data for neutral Xe@C_{60} with experimental data for charged Xe@C_{60}^{+} [3] is appropriate because [1, 27] charging the C_{60} shell positively does nothing to the photoionization cross section (as a function of photon energy) except to increase the threshold energy.

Results for depicted in Fig. 3 $\beta_{4d}(\omega)$ of Xe@C_{60} illustrate the impacts of the confining square-well and diffuse potentials (when compared with free Xe data) as well as electron correlation (when RPAE is compared with HF data) on the $\beta_{4d}(\omega)$ spectrum. They are self-explanatory. We omit commenting on them.

To conclude the discussion on the Xe@C_{60} $4d$ photoionization we note that very recently [19] new theoretical data on the Xe@C_{60} $4d$ photoionization were reported. Although being obtained on the basis of exactly the same approximations as in the works of three other independent theoretical groups [13, 15, 28] [i.e., RPAE (or relativistic RPAE in Ref. [15]) & square-well-potential with identical parameters) they differ strongly from results of all the three cited theoretical groups (the latter are in the mutual agreement with each other). We conclude that something was done wrong in work [19], most likely because of a very peculiar way those authors chose for solving HF equations in the presence of a square-well potential confinement.

2. The $5p$ photoionization of Xe@C_{60}

Commenting on $\sigma_{5p}(\omega)$ of Xe@C_{60}, Fig. 4(a), we note that a very strong maximum at the $5p$ threshold as well as a much less developed oscillation at about 30 eV carry the confinement resonance origin. Interesting, they are absent in $\beta_{5p}(\omega)$.

As for $\sigma_{5p}(\omega)$ beyond of a 100 eV energy, it dominates by the three strong resonances. Their positions approximately match the positions of confinement resonances in $\sigma_{4d}(\omega)$, Fig. 3. Exactly the same situation has recently [29] been predicted and explained for the Xe@C_{60} $5p$ generalized oscillator strength in a fast electron impact ionization of Xe@C_{60}. As in Ref. [29], the present study reveals that the resonances in $\sigma_{5p}(\omega)$ beyond 100 eV are induced by the confinement resonances in the $4d$ photoionization channels, via interchannel coupling. Thus, the resonances in $\sigma_{5p}(\omega)$ beyond 100 eV are of the *correlation* confinement resonance nature. Correlation confinement resonances in A@C_{60} photoionization [1, 4, 13, 15, 30] are resonances that emerge in the photoionization of an outer subshell due to interference of transitions from the outer subshell ($5p$ in our case) with confinement resonances emerging in *inner* shell transitions ($4d$ transitions in our case), via interchannel coupling. These correlation confinement resonances in the Xe@C_{60} $5p$ spectra are strong because of, as in free Xe, a strong intershell correlation coupling between $4d$ on $5p$ transitions.

Furthermore, one can see from Fig. 4(b) that correlation confinement resonances show up prominently in $\beta_{5p}(\omega)$ of Xe@C_{60} as well [resonances at about 85 and 100 eV in $\beta_{5p}(\omega)$]. This is another interesting prediction of the present study.

Finally, we note, that we omitted the C_{60} dynamical polarization impact [10, 31] (and references therein) on the Xe@C_{60} $5p$ ionization which is known to be significant up to about 70 eV of photon energy. The omission is justified because the primarily aim of the present paper is to study differences between effects of the confining square-well-potential and diffuse potential on A@C_{60} photoionization, and the C_{60} dynamical polarization cannot alter said differences.

[1] V. K. Dolmatov, in *Theory of Confined Quantum Sytems: Part Two*, edited by J. R. Sabin and E. Brändas, Advances in Quantum Chemistry (Academic Press, New York, 2009), Vol. **58**, pp. 13-68.

[2] V. K. Dolmatov, J. P. Connerade, A. S. Baltenkov, and S. T. Manson, Radiat. Phys. Chem. **70**, 417 (2004).

[3] A. L. D. Kilcoyne, A. Aguilar, A. Müller, S. Schippers, C. Cisneros, G. Alna'Washi, N. B. Aryal, K.K. Baral, D. A. Esteves, C. M. Thomas, and R. A. Phaneuf, Phys. Rev. Lett. **105**, 213001 (2010).

[4] M. E. Madjet, T. Renger, D. E. Hopper, M.A. McCune, H. S. Chakraborty, J.-M. Rost, and S. T. Manson, Phys. Rev. A **81**, 013202 (2010).

[5] M. Ya. Amusia, A. S. Baltenkov, L. V. Chernysheva, Z. Felfi, and A. Z. Msezane, J. Phys. B **38**, L169 (2005).

[6] G. Yu. Kashenock and R. H. Pratt, J.Phys.: Conf. Ser. **194**, 022055 (2009).

[7] M. J. Puska and R. M. Nieminen, Phys. Rev. A **47**, 1181 (1993).

[8] Y. B. Xu, M. Q. Tan, and U. Becker, Phys. Rev. Lett. **76**, 3538 (1996).

[9] W. Even, J. Smith, M. W. Roth, and H. A. Schuessler, Int. J. Mol. Sci. **5**, 333 (2004).

[10] S. Lo, A. V. Korol, and A. V. Solov'yov, J. Phys. B **40** (2007) 3973.

[11] M. Neek-Amal, G. Tayebirad, and R. Asgari, J. Phys. B **40** (2007) 1509.

[12] S. A. Ndengué and O. Motapon, J. Phys. B **41**, 045001 (2008).

[13] V. K. Dolmatov and S. T. Manson, J. Phys. B **41**, 165001 (2008).

[14] D. M. Mitnik, J. Randazzo, and G. Gasaneo, Phys. Rev. A **78**, 062501 (2008).

[15] Karan Govil, A. J. Siji, and P. C. Deshmukh, J. Phys. B **42** 065004 (2009).

[16] J. A. Ludlow, T. G. Lee, and M. S. Pindzola, Phys. Rev. A **81**, 023407 (2010).

[17] F. D. Colavecchia, G. Gasaneo, and D. Mitnik, J. At. Mol. Opt. Phys. **2011**, 817034 (2011).

[18] A. N. Grum-Grzhimailo, E. V. Gryzlova, and S. I. Strakhova, J. Phys. B **44**, 235005 (2011).

[19] Z. Chen and A. Msezane, Eur. Phys. J. D **65**, 353 (2011).

[20] O. Frank and J.-M. Rost, Chem. Phys. Lett. **271**, 367 (1997).

[21] G. Wendin and B. Wästberg, Phys. Rev. B **48**, 14764 (1993).

[22] E. M. Nascimento, F. V. Prudente, M. N. Guimarães, and A. M. Maniero, J. Phys. B **44**, 015003 (2011).

[23] A. B. Patel and H. S. Chakraborty, J. Phys. B **44**, 191001 (2011).

[24] M. Ya. Amusia and L. V. Chernysheva, *Computation of Atomic Processes* (IOP Publishing Ltd., Bristol, 1997).

[25] J. W. Cooper, Phys. Rev. A **47**, 1841 (1993).

[26] V. K. Dolmatov and D. A. Keating, J. Phys.: Conf. Ser. (in print); e-print arXiv:1109.5292v1 (2011).

[27] V. K. Dolmatov and S. T. Manson, Phys. Rev. A **73**, 013201, (2006).

[28] M. Ya. Amusia and L. V. Chernysheva, private communication (2011).

[29] M. Ya. Amusia, L. V. Chernysheva, and V. K. Dolmatov, Phys. Rev. A **84**, 063201 (2011).

[30] V. K. Dolmatov, G. T. Craven, E. Guler, and D. Keating, Phys. Rev. A **80**, 035401 (2009).

[31] M. Ya. Amusia, A. S. Baltenkov, and L. V. Chernysheva, JETP Lett. **87**, 200 (2008).